

# Research Paper

## Chester Treadmill Police Tests as Alternatives to 15-metre Shuttle Running

**Dr Michael Morris**, Department of Clinical Sciences & Nutrition, University of Chester, Chester, UK

**Elizabeth Parker**, Department of Clinical Sciences & Nutrition, University of Chester, Parkgate Road, Chester, CH1 4BJ

**Professor Kevin Sykes**, Emeritus Professor of Occupational Health and Workplace Fitness, University of Chester

### Abstract

**Background:** Police Officers require a specific level of aerobic fitness to allow them to complete Personal Safety Training (PST) and specialist roles. Officers' aerobic fitness is assessed using the 15m multi-stage fitness test, however, due to the agility required and risk of injury, two alternative treadmill tests have been designed to predict four of the key minimum  $\dot{V}O_2$  criteria of 35, 41, 46 and 51 ml.kg<sup>-1</sup>.min<sup>-1</sup>.

**Aims:** To investigate the validity and reliability of Chester Treadmill Police Walk Test and Chester Treadmill Police Run Test.

**Methods:** 78 UK Police officers (18 = female) completed the CTPWT (n=53) or CTPRT (n=35), or both; generating a total of 88 data sets. To assess reliability 43 participants returned for a second visit (T2), to repeat the treadmill test.

**Results:** Mean differences between predicted and actual  $\dot{V}O_2$  at 35, 41, 46 and 51 ml.kg<sup>-1</sup>.min<sup>-1</sup> were as follows -1.1, -2.1, -0.1 and -1.2ml.kg<sup>-1</sup>.min<sup>-1</sup>. Despite a significant under prediction (p=.001) a minimum of 92% of participants were within 10% of target  $\dot{V}O_2$  at all levels. There was no significant difference between actual and predicted  $\dot{V}O_2$  in the CTPRT, at 46 ml.kg<sup>-1</sup>.min<sup>-1</sup> (T1 46.0 ± 1.4 or T2 45.1 ± 1.3 ml.kg<sup>-1</sup>.min<sup>-1</sup>). Similarly, there was no significant difference at 51 ml.kg<sup>-1</sup>.min<sup>-1</sup> (T2 50.5 ± 1.4 ml.kg<sup>-1</sup>.min<sup>-1</sup>). We observed no differences for gender or trial. 95% Limits of Agreement were at worst T1-T2 -0.25 ± 4.0ml.kg<sup>-1</sup>.min<sup>-1</sup>.

**Conclusions:** The CTPWT and the CTPRT provide a valid and reliable alternative to the 15m MSFT.

**Keywords:** occupational, fitness, exercise testing, Police, fitness standards, predictive, treadmill test

## Introduction

Maximal oxygen consumption ( $\dot{V}O_{2max}$ ) is a commonly used measure of aerobic, or cardiorespiratory, fitness<sup>1</sup>. This measure has been associated with cardiometabolic health, mortality and athletic performance alike<sup>2-6</sup>. Whilst it is desirable to measure  $\dot{V}O_{2max}$  using online gas analysis, alternative tests have been developed to allow for the prediction of  $\dot{V}O_{2max}$  without the use of specialised equipment or ergometry<sup>7,8</sup>. The use of these submaximal exercise tests to predict  $\dot{V}O_{2max}$  has been widely investigated by researchers. Although there are noted limitations of submaximal exercise testing such as the reliance upon an accurate HR, power output and  $\dot{V}O_2$  relationship there are many submaximal protocols which are accepted to be an effective means of assessing cardiovascular fitness<sup>9,10</sup>.

Similar to military personnel and firefighters<sup>11</sup>, police officers are required to attend regular fitness testing to continue in operation. Following the Winsor report and based upon research carried out by Brewer<sup>12</sup> the College of Policing set out required standards of aerobic fitness for all operational officers (Table 1). To achieve the fitness required for personal safety training (PST) a  $\dot{V}O_{2max}$  of at least  $35\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  is necessary<sup>12</sup>, with increasing requirements for specialist roles (Table 1). These requirements are assessed using a 15m multi-stage fitness test (MSFT) which has been validated against the 20m MSFT in an unpublished study<sup>13</sup>. This 15m MSFT has also been compared to PST and specialist roles physiological demands<sup>14</sup> and to  $\dot{V}O_{2max}$  in prior research<sup>15</sup>. However, the validity and reliability of shuttle testing to predict  $\dot{V}O_{2max}$  has been questioned among specific populations<sup>1,16</sup>. Prior to participation all officers are required to complete a medical questionnaire to ensure there are no medical contraindications such as musculoskeletal issues which may be aggravated by the twisting and turning associated with the MSFT. For this reason, some Forces have implemented an alternative fitness test (Chester Step Test, Astrand Cycle Test, and Chester Treadmill Walk Test), however the validity and reliability of the results may be questioned due to the lack of research investigating this. This paper will investigate the validity and reliability of Chester Treadmill Police Walk Test<sup>17</sup> and Chester Treadmill Police Run Test<sup>18</sup> in predicting four of the key  $\dot{V}O_2$  values laid out by Brewer<sup>12</sup> highlighted in Table 1. These treadmill tests were developed by Sykes (2015) to predict specified  $\dot{V}O_2$  values (Table 1) using well established ACSM metabolic equations<sup>19</sup>.

Table 1: Police Fitness Standards (Adapted from Brewer <sup>12</sup>)

<b>Unit</b>	<b>Recommended Standard (Level : Shuttle)</b>	<b>Est. Aerobic Capacity* (ml.kg<sup>-1</sup>.min<sup>-1</sup>)**</b>
<b>Personal Safety Training</b>	<b>5 : 4</b>	<b>35</b>
<b>Marine Police Unit</b>	<b>5 : 4</b>	<b>35</b>
<b>CBRN</b>	<b>5 : 4</b>	<b>35</b>
<b>Method of Entry</b>	<b>5 : 4</b>	<b>35</b>
Dog Handler	5 : 7	36
Mounted Branch	5 : 7	36
Police Cyclist	5 : 8	36
Police Support Unit	6 : 3	37
Air Support	6 : 4	37
Police Divers	6 : 8	39
Marine Police (Tactical Skills)	7 : 2	40
<b>Authorised Firearms Officer</b>	<b>7 : 6</b>	<b>41</b>
<b>Armed Response Vehicle</b>	<b>9 : 4</b>	<b>46</b>
<b>Dynamic Intervention AFO</b>	<b>10: 5</b>	<b>51</b>

\* Aerobic Capacity must be at least this value in order to attain the Shuttle standard

\*\* ml.kg<sup>-1</sup>.min<sup>-1</sup> values rounded to nearest whole number

## Methods

A total of 78 UK Police officers (18 = female) volunteered to take part in the study (age:  $42 \pm 7$  years; Height:  $1.8 \pm 0.1$ m; Weight:  $82.1 \pm 15.2$ kg; BMI:  $26.1 \pm 3.5$ ). The study aimed to recruit 40 participants per group (CTPWT and CTPRT) as per recommendations by Atkinson and Nevill<sup>20</sup>. All participants completed written, informed consent and health screening prior to taking part in the research project, which gained ethical approval from the University of Chester. Participants attended the University of Chester on two separate occasions having abstained from caffeine, alcohol and vigorous exercise for 24 hours. Blood pressure and resting heart rate were recorded (Omron, Germany) along with body mass (kg) and stature (cm) (Seca, Germany). During each testing day participants took part in the relevant treadmill test according to their self-reported performance of the 15m MSFT, (i.e. which level they are able to run to during the MSFT; 5:4, 7:6, 9:4 or 10:5) which participants were all familiar with. All participants were also familiar with treadmill walking and/or running prior to attending the University. Participants had a minimum of 24 hours between testing days to allow adequate recovery time and repeatability was assessed within two weeks to limit the effect of time or changes to fitness. During exercise, heart rate (HR) (Polar, Finland), RPE<sup>21</sup> and oxygen consumption ( $\dot{V}O_2$ ) were measured. Oxygen consumption was measured with online gas analysis (Metamax 3B, Cortex, Germany) and data was averaged to 10 seconds for subsequent analysis. Whilst there are limitations associated with the use of time average data smoothing,<sup>22</sup> 10s averaging was employed to allow for future studies to directly compare with 15m MSFT  $\dot{V}O_2$  data, for which 10s averaging would be necessary to discern between shuttle levels. The use of this time-second averaging method is also supported by its use in similar studies<sup>23</sup>. Observation of the data by researchers showed no difference between the final two 10-s average periods recorded per stage.  $\dot{V}O_2$  values for the CTPWT and CTPRT (Sykes, 2015) were predicted using well established ACSM equations<sup>19</sup>.

Data was analysed using SPSS for Windows (Version 22) and alpha was set at the .05 level. Normality of data was checked using Shapiro-Wilk statistic and descriptive statistics (mean  $\pm$  SD) were computed. To investigate the difference between actual and predicted  $\dot{V}O_2$  values (validity) one sample t-tests were applied and independent t-tests to compare gender differences. The test-retest differences (reliability) were investigated using paired sample t-tests, 95% limits of agreement (LoA) (bias  $\pm$   $1.96 \times SD_{diff}$ ), Bland Altman LoA, typical error and intra-class correlation coefficient.

Table 2 Chester Treadmill Police Walk Test and Chester Treadmill Police Run Test Protocols <sup>17 18</sup> )

<b>Chester Treadmill Police Walk Test</b>			
ACSM equation for walking: $\dot{V}O_2$ (ml.kg-1.min-1) = (0.1 . S) + (1.8 . S . G) + 3.5 ml.kg-1.min-1 <sup>24</sup>			
<b>Level</b>	<b>Time (mins)</b>	<b>Treadmill Gradient</b>	<b>Predicted O<sub>2</sub> cost (ml.kg-1.min-1)*</b>
Speed: 6.0km/hr			
1	0-2	0%	14
2	2-4	3%	19
3	4-6	6%	24
4	6-8	9%	30
5	8-10	12%	35
6	10-12	15%	41
<b>Chester Treadmill Police Run Test</b>			
<b>Level</b>	<b>Time (mins)</b>	<b>Treadmill Gradient</b>	<b>Predicted O<sub>2</sub> cost (ml.kg-1.min-1)*</b>
Speed: 10.4km/hr			
ACSM equation for running: $\dot{V}O_2$ (ml.kg-1.min-1) = (0.2 . S) + (0.9 . S . G) + 3.5 ml.kg-1.min-1 <sup>24</sup>			
1	0-2	0%	38
2	2-4	2%	41
3	4-6	4%	44
4	6-8	5%	46
5	8-10	8%	51

S = speed in m.min-1; G = percent grade expressed as a fraction

\*ml.kg-1.min-1 values rounded to nearest whole number

## Results

Fifty three participants (15 female) completed up to Level 5 (35ml.kg<sup>-1</sup>.min<sup>-1</sup>) on CTPWT once, with 30 (9 female) completing to this level twice, whilst 52 participants (15 female) completed up to Level 6 (41ml.kg<sup>-1</sup>.min<sup>-1</sup>) on CTPWT once, with 28 (8 female) completing to this level twice.

Thirty five participants (9 female) completed up to Level 4 (46ml.kg<sup>-1</sup>.min<sup>-1</sup>) on CTPRT with 13 (1 female) completing to this level twice, whilst 34 (1 female) participants completed to Level 5 (51ml.kg<sup>-1</sup>.min<sup>-1</sup>) on CTPRT once, with 13 (1 female) completing to this level twice.

*Table 3 Validity CTPWT and CTPRT*

<b>Treadmill Level (Time)</b>	<b>Predicted <math>\dot{V}O_2</math> (ml.kg<sup>-1</sup>.min<sup>-1</sup>)</b>	<b>Trial 1 Actual <math>\dot{V}O_2</math> (ml.kg<sup>-1</sup>.min<sup>-1</sup>) (95% Confidence Interval)</b>	<b>Trial 2 Actual <math>\dot{V}O_2</math> (ml.kg<sup>-1</sup>.min<sup>-1</sup>) (95% Confidence Interval)</b>
<b>Treadmill Walk test</b>			
Level 5 (10 mins)	35	34.0 ± 1.8* (33.5 – 34.7)	34.3 ± 1.8* (33.7 – 35.0)
Level 6 (12 mins)	41	39.0 ± 2.3* (38.4 – 40.0)	39.0 ± 2.5* (38.0 – 40.0)
<b>Treadmill Run Test</b>			
Level 4 (8 mins)	46	45.9 ± .7 (45.6 – 46.5)	46.0 ± 1.4 (45.1 – 46.7)
Level 5 (10 mins)	51	50.6 ± 1.2* (49.9 – 51.3)	50.5 ± 1.4 (49.6 – 51.4)

\*denotes significant difference from predicted  $\dot{V}O_2$  p < .05

### **Chester Treadmill Walk Test**

In trial 1 of the CTPWT there was a significant under prediction between recorded  $\dot{V}O_2$  and predicted  $\dot{V}O_2$  at 35 ml.kg<sup>-1</sup>.min<sup>-1</sup> (34.0 ± 1.8 ml.kg<sup>-1</sup>.min<sup>-1</sup>; p = .001) and 41 ml.kg<sup>-1</sup>.min<sup>-1</sup> (39.0 ± 2.3 ml.kg<sup>-1</sup>.min<sup>-1</sup>; p = .001) and in trial 2 at 35 ml.kg<sup>-1</sup>.min<sup>-1</sup> (34.3 ± 1.8 ml.kg<sup>-1</sup>.min<sup>-1</sup>; p = .049) and 41 ml.kg<sup>-1</sup>.min<sup>-1</sup> (39.0 ± 2.5 ml.kg<sup>-1</sup>.min<sup>-1</sup>; p = .001). Despite this, 92% of participants (100% of females) in trial 1 and 93% of participants (100% of females) in trial 2 at Level 5 (minimum requirement for officers undertaking PST) were within 10% (i.e. equivalent to 1MET = 3.5 ml.kg<sup>-1</sup>.min<sup>-1</sup>) of the target  $\dot{V}O_2$  value of 35 ml.kg<sup>-1</sup>.min<sup>-1</sup>. All participants in trial 1 and 82% of participants in trial 2 (94% of females) at Level 6 were within 10% (4.1 ml.kg<sup>-1</sup>.min<sup>-1</sup>) of predicted values. There was no significant difference between males and females in trial 1 at 35 ml.kg<sup>-1</sup>.min<sup>-1</sup> (p = .628), trial 2 at 35 ml.kg<sup>-1</sup>.min<sup>-1</sup> (0.76), trial 1 at 41 ml.kg<sup>-1</sup>.min<sup>-1</sup> (p=.88) or trial 2 at 41 ml.kg<sup>-1</sup>.min<sup>-1</sup> (p=.9).

The test-retest differences (reliability) of the CTPWT are summarised in Table 4, with comparisons across two trials. Paired t-tests revealed no significant differences between Trial 1 and Trial 2 at any level. 95% Limits of Agreement between Trial 1 and Trial 2 were as follows for 35 and 41 ml.kg<sup>-1</sup>.min<sup>-1</sup> respectively; -0.25 ± 4.0 ml.kg<sup>-1</sup>.min<sup>-1</sup>, 0.15 ± 2.8 ml.kg<sup>-1</sup>.min<sup>-1</sup>. Bland-Altman plots (Figures 1-2) showed acceptable limits of agreement (LoA) between Trial 1 and Trial 2.

### **Chester Treadmill Police Run Test**

Actual  $\dot{V}O_2$  and predicted  $\dot{V}O_2$  were significantly different (p<.05) at 51 ml.kg<sup>-1</sup>.min<sup>-1</sup> in trial 1 of the CTPRT (50.6 ± 1.2 ml.kg<sup>-1</sup>.min<sup>-1</sup>). Despite this, 92% of participants in trial 1 (100% of females) and all participants in trial 2 at Level 4 were within 10% (4.6 ml.kg<sup>-1</sup>.min<sup>-1</sup>) of predicted values. 94% of participants (88% of females) in trial 1 and 100% in trial 2 at Level 5 were within 10% (5.1 ml.kg<sup>-1</sup>.min<sup>-1</sup>) of predicted values. There was no significant difference between actual and predicted  $\dot{V}O_2$  in the CTPRT, at 46 ml.kg<sup>-1</sup>.min<sup>-1</sup> during trial 1 (46.0 ± 1.4 ml.kg<sup>-1</sup>.min<sup>-1</sup>) or trial 2 (45.1 ± 1.3 ml.kg<sup>-1</sup>.min<sup>-1</sup>). Similarly, there was no significant difference at 51 ml.kg<sup>-1</sup>.min<sup>-1</sup> during trial 2 (50.5 ± 1.4 ml.kg<sup>-1</sup>.min<sup>-1</sup>). There was no significant difference between males and females in trial 1 at 46 ml.kg<sup>-1</sup>.min<sup>-1</sup> (p = .9), trial 2 at 46 ml.kg<sup>-1</sup>.min<sup>-1</sup> (p = .6), trial 1 at 51 ml.kg<sup>-1</sup>.min<sup>-1</sup> (p = .4) or trial 2 at 51 ml.kg<sup>-1</sup>.min<sup>-1</sup> (p = .7).

The test-retest differences (reliability) of the CTPWT and CTPRT are summarised in Table 4, with comparisons across two trials. Paired t-tests revealed no significant differences between Trial 1 and Trial 2 at any level. 95% Limits of Agreement between Trial 1 and Trial 2 were as follows for 46 and 51 ml.kg<sup>-1</sup>.min<sup>-1</sup> respectively; 0.17 ± 2.8 ml.kg<sup>-1</sup>.min<sup>-1</sup>, 0.08 ± 2.3 ml.kg<sup>-1</sup>.min<sup>-1</sup>. Bland-Altman plots (Figures 3-4) showed acceptable limits of agreement (LoA) between Trial 1 and Trial 2.

Table 4 Test-retest differences of CTPWT and CTPRT across two repeated trials.

Treadmill Level (Time)	95% LoA <sup>1</sup> (bias ± 1.96 x SD <sub>diff</sub> )	ICC	Typical error <sup>1</sup>
<b>Chester Treadmill Walk test (CTPWT)</b>			
Level 5 (10 mins)	-0.25 ± 4.0	.37	± 1.33
Level 6 (12 mins)	0.15 ± 2.8	.89	± .39
<b>Chester Treadmill Run Test (CTPRT)</b>			
Level 4 (8 mins)	0.17 ± 2.8	.28	± 1.66
Level 5 (10 mins)	0.08 ± 2.3	.80	± 0.51

<sup>1</sup>ml·kg<sup>-1</sup>·min<sup>-1</sup>




Figure 1 Bland Altman Plots of reliability between trials



## Discussion

Whilst our results show a statistically significant difference at some levels of the predictive treadmill tests, the magnitude of these differences are likely negligible in a practical setting with all mean differences between .06 and 2 ml·kg<sup>-1</sup>·min<sup>-1</sup>. Previous research examining treadmill protocols to predict  $\dot{V}O_{2max}$  have reported error of between 11-18% and deem these levels to be unacceptable<sup>23</sup><sup>25</sup><sup>26</sup>. Drew-Nord, et al.<sup>27</sup> report over-estimations of between 1-2 METs with two predictive treadmill protocols, which would equate to around 3.5-7ml·kg<sup>-1</sup>·min<sup>-1</sup> similar to findings by Zwiren, et al.<sup>9</sup> and Tierney, et al.<sup>28</sup> who report SEE of between 2.9 and 5.2 ml·kg<sup>-1</sup>·min<sup>-1</sup>. This research reports variability much greater than shown within our study yet accepts tests as suitable for use, thus further supporting the findings of our study. Some evidence suggests that prediction equations overestimate the fitness of lower fit individuals and under predict the fitness of higher fit individuals<sup>23</sup><sup>26</sup>, however our study did not investigate aerobic capacity of participants thus we are unable to comment on this. Confidence intervals of all levels do show a slight underestimation of  $\dot{V}O_2$  (table 1) however the magnitude of this is very low and arguably negligible in the practical setting. From the findings of this study, specialist units, who have specific  $\dot{V}O_2$  values to achieve, have been set individual time targets on the CTPWT (see Appendix, Table 6 and 7).

A limitation of the current study is the reliance upon tables provided by Brewer<sup>12</sup> (Table 1) which unfortunately gives no reference to confidence levels, or error margins, of estimated aerobic capacity. These values were devised by the equations provided by Roehampton University of Surrey<sup>15</sup> which similarly gave no indication of confidence levels. Therefore, the scarcity of information makes comparison of results somewhat difficult. However, research into the 20m MSFT provides 95% LoA of ~6ml<sup>16</sup>, suggesting SD of ~3ml, greater than ours (average SD of 1.6 ml·kg<sup>-1</sup>·min<sup>-1</sup>). It could be suggested that these SD and LoA provided by Aandstad, et al.<sup>16</sup> would be similar to Roehampton data, which would strengthen the findings and the predictive value of our tests to being able to predict  $\dot{V}O_2$  within these margins.

A particular strength of this study is that all participants were serving UK police officers, thus the findings can be generalized to the relevant population. Furthermore, the study included both male and female participants which prior similar research has commonly failed to address<sup>23</sup><sup>26</sup><sup>27</sup>. Interestingly, our analysis showed there was no gender bias for either the CTPWT or the CTPRT. Although great efforts were made, we did not have any black or minority ethnicity (BME) participants volunteer to take part in the study which weakens the generalizability of the findings. There are also accepted errors innate in the use of metabolic analysers<sup>29</sup> however the unit used within this study has been reported to be stable and reliable<sup>30</sup> and was used consistently among all trials to control for variation between units. Due to operational constraints not all officers were able

to return for trial two testing, thus decreasing sample size of the reliability study, however, analysis of the data demonstrated excellent reliability across trials. Using LoA statistics, the CTPWT and CTPRT may over or under-predict  $\dot{V}O_2$  values by between 2-4 ml·kg<sup>-1</sup>·min<sup>-1</sup>. Typical error values between 0.51ml·kg<sup>-1</sup>·min<sup>-1</sup> and 1.66 ml·kg<sup>-1</sup>·min<sup>-1</sup> further support the strength of the tests repeatability.

To conclude, whilst the statistical analysis of our study suggests there are statistical differences between predicted  $\dot{V}O_2$  values and actual  $\dot{V}O_2$  values, these variations are thought to be negligible within the practical setting and are much closer to predicted values than previous research has reported. Reliability data shows that there is no need for a familiarisation test for officers using either the CTPWT or CTPRT. Both the CTPWT and the CTPRT are therefore deemed as suitable alternative tests to the 15m MSFT for any officer undertaking PST and for various specialist units (Table 6 and 7).

### Key Points

- The CTPWT and CTPRT are valid tests to predict the  $\dot{V}O_2$  values set out by Brewer (2010).
- The CTPWT slightly under predicted  $\dot{V}O_2$  however the practical relevance is negligible and has minimal impact on classification of participants in attaining the required  $\dot{V}O_2$  for their roles.
- The CTPWT and CTPRT are reliable tests and a familiarisation test isn't deemed necessary in the practical setting.

### Funding

Funding for the research project was provided by Cumbria Constabulary.

### Conflicts of Interest

The authors declare no conflicts of interest.

### Acknowledgements

We would like to acknowledge the invaluable support of numerous Constabularies and to all Police Officers who volunteered their time to participate in the project. We would also like to thank Paul Buckle and Paul Telford for their expertise, support and knowledge throughout the project.

## References

1. Grant S, Corbett K, Amjad AM, et al. A comparison of methods of predicting maximum oxygen uptake. *British journal of sports medicine* 1995;29(3):147-52. [published Online First: 1995/09/01]
2. Cosgrove MJ, Wilson J, Watt D, et al. The relationship between selected physiological variables of rowers and rowing performance as determined by a 2000 m ergometer test. *Journal of sports sciences* 1999;17(11):845-52. doi: 10.1080/026404199365407 [published Online First: 1999/12/10]
3. Lee SJ, Kuk JL, Katzmarzyk PT, et al. Cardiorespiratory Fitness Attenuates Metabolic Risk Independent of Abdominal Subcutaneous and Visceral Fat in Men. *Diabetes Care* 2005;28:6.
4. Earnest CPA, E. G. ; Xuemei, S.; Duck-chul, L.; Church, T. S.; Blair, S. N. Maximal Estimated Cardiorespiratory Fitness, Cardiometabolic Risk Factors, and Metabolic Syndrome in the Aerobic Centre Longitudinal Study. *Mayo Clinic Proceedings* 2013;88(3):259-70.
5. Shuval K, Finley CE, Barlow CE, et al. Sedentary behavior, cardiorespiratory fitness, physical activity, and cardiometabolic risk in men: the cooper center longitudinal study. *Mayo Clinic Proceedings* 2014;89(8):1052-62. doi: 10.1016/j.mayocp.2014.04.026
6. Wiswell RA, Jaque SV, Marcell TJ, et al. Maximal aerobic power, lactate threshold, and running performance in master athletes. *Med Sci Sports Exerc* 2000;32(6):1165-70. [published Online First: 2000/06/22]
7. Sykes K, Roberts A. The Chester step test—a simple yet effective tool for the prediction of aerobic capacity. *Physiotherapy* 2004;90(4):183-88. doi: <http://dx.doi.org/10.1016/j.physio.2004.03.008>
8. Siconolfi SF, Cullinane EM, Carleton RA, et al. Assessing VO<sub>2</sub>max in epidemiologic studies: modification of the Astrand-Rhyming test. *Medicine and science in sports and exercise* 1982;14(5):335-38.
9. Zwiren LD, Freedson PS, Ward A, et al. Estimation of VO<sub>2</sub>max: a comparative analysis of five exercise tests. *Res Q Exerc Sport* 1991;62(1):73-8. doi: 10.1080/02701367.1991.10607521 [published Online First: 1991/03/01]
10. Bennett H, Parfitt G, Davison K, et al. Validity of Submaximal Step Tests to Estimate Maximal Oxygen Uptake in Healthy Adults. *Sports medicine (Auckland, NZ)* 2016;46(5):737-50. doi: 10.1007/s40279-015-0445-1 [published Online First: 2015/12/17]
11. Sykes K. Fitness for Fire and Rescue. Standards, Protocols and Policy. In: Stevenson R, Wilsher P, eds. United Kingdom: Firefit Steering Group, 2009.
12. Brewer J. Job Related Fitness Tests for Police Officer Specialist Posts. United Kingdom: National Police Improvement Agency, 2010.
13. Loughborough University of Technology. Recommendations For The Standards Of Fitness Of Metropolitan Police Officers. United Kingdom: Loughborough University of Technology, 1995.
14. Brewer J, Buckle P, Castle P. The Use of Work Place Physiological Measurements to Establish the Minimum Fitness Standards Required for Entry into the United Kingdom Police Service. *Athletic Enhancement* 2014;2013
15. Roehampton University of Surrey. Report on the Validation of the 15m Work Related Fitness Test. United Kingdom: Roehampton University of Surrey, 2003.
16. Aandstad A, Holme I, Berntsen S, et al. Validity and reliability of the 20 meter shuttle run test in military personnel. *Mil Med* 2011;176(5):513-8. [published Online First: 2011/06/04]
17. Sykes K. The Chester Treadmill Police Walk Test. United Kingdom: Cartwright Fitness, 2015.
18. Sykes K. The Chester Treadmill Police Run Test. United Kingdom: Cartwright Fitness, 2015.
19. American College of Sports Medicine. Advanced Fitness Assessment and Exercise Prescription. 7th ed. Leeds, United Kingdom: Human Kinetics 2013.
20. Atkinson G, Nevill AM. Statistical methods for assessing measurement error (reliability) in variables relevant to sports medicine. *Sports medicine (Auckland, NZ)* 1998;26(4):217-38. [published Online First: 1998/11/20]

21. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982;14(5):377-81. [published Online First: 1982/01/01]
22. Robergs RA, Dwyer D, Astorino T. Recommendations for Improved Data Processing from Expired Gas Analysis Indirect Calorimetry. *Sports Medicine* 2010;40(2):95-111. doi: <http://dx.doi.org/10.2165/11319670-000000000-00000>
23. Dolezal BA, Barr D, Boland DM, et al. Validation of the firefighter WFI treadmill protocol for predicting VO<sub>2</sub> max. *Occupational medicine (Oxford, England)* 2015;65(2):143-6. doi: 10.1093/occmed/kqu189 [published Online First: 2015/01/09]
24. Medicine ACoS. Metabolic Equations for Gross VO<sub>2</sub> in Metric Units 2015 [accessed 13/09/2016 2016.
25. Mier CM, Gibson AL. Evaluation of a treadmill test for predicting the aerobic capacity of firefighters. *Occupational medicine (Oxford, England)* 2004;54(6):373-8. doi: 10.1093/occmed/kqh008 [published Online First: 2004/09/07]
26. Klaren RE, Horn GP, Fernhall B, et al. Accuracy of the VO<sub>2</sub>peak prediction equation in firefighters. *Journal of occupational medicine and toxicology (London, England)* 2014;9:17. doi: 10.1186/1745-6673-9-17 [published Online First: 2014/05/27]
27. Drew-Nord DC, Myers J, Nord SR, et al. Accuracy of peak VO<sub>2</sub> assessments in career firefighters. *Journal of occupational medicine and toxicology (London, England)* 2011;6(1):25. doi: 10.1186/1745-6673-6-25 [published Online First: 2011/09/29]
28. Tierney MT, Lenar D, Stanforth PR, et al. Prediction of aerobic capacity in firefighters using submaximal treadmill and stairmill protocols. *Journal of strength and conditioning research / National Strength & Conditioning Association* 2010;24(3):757-64. doi: 10.1519/JSC.0b013e3181c7c282 [published Online First: 2010/02/11]
29. Hodges LD, Brodie DA, Bromley PD. Validity and reliability of selected commercially available metabolic analyzer systems. *Scandinavian journal of medicine & science in sports* 2005;15(5):271-9. doi: 10.1111/j.1600-0838.2005.00477.x [published Online First: 2005/09/27]
30. Macfarlane DJ, Wong P. Validity, reliability and stability of the portable Cortex Metamax 3B gas analysis system. *European journal of applied physiology* 2012;112(7):2539-47. doi: <http://dx.doi.org/10.1007/s00421-011-2230-7>

## Appendix

*Table 3 CTPWT target time.*

Unit	Recommended Standard (Level : Shuttle)	Est. Aerobic Capacity* (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )**	CTPWT Target Time (minutes:seconds)*
Personal Safety Training	5 : 4	35	10:00
Marine Police Unit	5 : 4	35	
CBRN	5 : 4	35	
Method of Entry	5 : 4	35	
Dog Handler	5 : 7	36	10:20
Mounted Branch	5 : 7	36	
Police Cyclist	5 : 8	36.	
Police Support Unit	6 : 3	37	10:40
Air Support	6 : 4	37	
Police Divers	6 : 8	39	11:20
Marine Police (Tactical Skills)	7 : 2	40	11:40
Authorised Firearms Officer	7 : 6	41	12:00

*Table 4 CTPRT target times*

Unit	Recommended Standard (Level : Shuttle)	Est. Aerobic Capacity* (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )**	CTPRT Target Time (minutes)
Armed Response Vehicle	9 : 4	46	10:00
Dynamic Intervention AFO	10 : 5	51	12:00

Table 5 Descriptive Statistics Trial 1 and Trial 2 CTPWT

Specialist Post	CTPWT Target Time (min:sec)	Est. Aerobic Capacity* (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )	T1 Actual $\dot{V}O_2$ (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )	T2 Actual $\dot{V}O_2$ (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )
Marine Police Unit	10:00	35	33.7 ± 1.8	33.6 ± 2.5
CBRN				
Method of Entry				
Dog Handler	10:20	36	34.9 ± 1.9	34.6 ± 3.3
Mounted Branch				
Police Cyclist				
Police Support Unit	10:40	37	35.6 ± 2.1	35.1 ± 3.0
Air Support				
Police Divers	11:20	39	37.3 ± 2.3	37.5 ± 3.4
Marine Police (Tactical Skills)	11:40	40	37.5 ± 2.8	38.0 ± 3.1
Authorised Firearms Officer	12:00	41	38.7 ± 2.1	39.1 ± 2.8

Table 6 Descriptive Statistics Trial 1 and Trial 2 CTPRT

Specialist Post	CTPRT Target Time (min:sec)	Est. Aerobic Capacity* (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )	T1 Actual $\dot{V}O_2$ (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )	T2 Actual $\dot{V}O_2$ (ml·kg <sup>-1</sup> ·min <sup>-1</sup> )
ARV	8:00	46	45.1 ± 1.9	45.5 ± 1.8
DIAFO	10:00	51	50.2 ± 1.8	50.1 ± 1.4




Figure 5: Mean VO<sub>2</sub> between 10 and 12 minutes of CTPWT and line of best fit using ACSM metabolic equations